# In the Supreme Court of the United States

PARENTS PROTECTING OUR CHILDREN, UA,

Petitioner,

v.

EAU CLAIRE AREA SCHOOL DISTRICT, WISCONSIN, ET AL.,

Respondents.

On Petition for a Writ of Certiorari to the United States Court of Appeals for the Seventh Circuit

# BRIEF OF THE MANHATTAN INSTITUTE AND DR. LEOR SAPIR AS AMICI CURIAE SUPPORTING PETITIONER

Ilya Shapiro
Counsel of Record
MANHATTAN INSTITUTE
52 Vanderbilt Ave.
New York, NY 10017
(212) 599-7000
ishapiro@manhattan.institute

July 8, 2024

# **QUESTION PRESENTED**

When a school district adopts an explicit policy to usurp parental decisionmaking authority over a major health-related decision—and to conceal this from the parents—do parents who are subject to such a policy have standing to challenge it?

# TABLE OF CONTENTS

QŢ	JES	STI	ON PRESENTED	. i
ТА	BL	ΕC	OF AUTHORITIES	iv
IN	TE	RES	ST OF AMICI CURIAE	1
BA			OUND AND SUMMARY OF JMENT	1
AR	RGU	JMI	ENT	3
I.	Social Transition Constitutes a Mental-Health Intervention for Children Who Would Otherwise Likely Desist in Their Adopted Gender Identity before Adulthood			
	A.	Th Int	edical Research Worldwide Demonstrates at Social Transition Is a Mental-Health tervention with Medical Implications for ildren and Adolescents	3
		1.	In most cases, childhood-onset gender dysphoria remits naturally by adulthood, but social transition may contribute to the persistence of gender dysphoria	5
		2.	Transgender identity in adolescents is also likely unstable.	9

B. Objections to the Persistence and Desistence Literature Do Not Withstand Scrutiny	13
C. Clinicians Have Not Demonstrated a Consistent Ability to Distinguish between Transgender and Gender-Nonconforming Youths	16
D. U.SBased Medical Groups Are Out of Step with World Health Authorities' Recognition of the Risks of Pediatric Social Transition	17
II. U.S. Courts Have Recognized Social Transition as a Medical Intervention	21
III.The Eau Claire Area School District's Gender Identity Support Policy Prevents Parents from Assessing the Benefits and Risks of Social Transition and Other Mental-Health Aspects of Gender Care	22
CONCLUSION	24

#### TABLE OF AUTHORITIES

# Cases Adams v. Sch. Bd. of St. Johns Cnty., Dodds v. U.S. Dep't of Educ., Doe v. Boyertown Area Sch. Dist., 897 F.3d 518 (3d Cir. 2018)......22 Grimm v. Gloucester Cnty. Sch. Bd., Lassiter v. Dep't of Soc. Servs., 452 U.S. 18 (1981)... 23 Parents for Privacy v. Dallas Sch. Dist. No. 2, Prince v. Massachusetts, 321 U.S. 158 (1943) ......... 23 Whitaker v. Kenosha Unified Sch., Dist., **Statutes** Other Authorities Am. Psych. Ass'n, DSM-5......14

Annelou L. C. de Vries & Peggy T. Cohen-Kettenis, Clinical Management of Gender Dysphoria in Children and Adolescents: The Dutch Approach, 59 J. Homosexuality 301 (2012)
Annelou L. C. de Vries et al., Reliability and Clinical Utility of Gender Identity-Related Diagnoses: Comparisons Between the ICD-11, ICD-10, DSM-IV, and DSM-5, 8 LGBT Health 133 (2021)
Annika Mutanen, A Professor Who Treats Adolescent Gender Anxiety Says No to Minors' Legal Gender Correction, Helsingin Sanomat, Jan. 27, 2023
Appendix A to Supplemental Expert Report of James Cantor, Ph.D., Boe v. Marshall, No. 2:22-cv-00184-LCB-CWB (M.D. Ala. June 24, 2024)
Benjamin Freedman, Equipoise and the Ethics of Clinical Research, 317 New Eng. J. Med. 141 (1987)
Christina M. Roberts et al., Continuation of Gender- Affirming Hormones Among Transgender Adolescents and Adults, 107 J. Clinical Endocrinology & Metabolism 3937 (2022)
E. Abruzzese et al., The Myth of "Reliable Research" in Pediatric Gender Medicine: A Critical Evaluation of the Dutch Studies— and Research that Has Followed, J. Sex & Marital Therapy 1 (2023)9-10, 11
Eli Coleman et al., Standards of Care for the Health of Transgender and Gender Diverse People, Version 8, 23 Int'l J. Transgender Health S1 (Suppl. 1) (2022)

Eli Coleman et al., Standards of Care for the Health of Transsexual, Transgender, and Gender Nonconforming People, Version 7, 13 Int'l J. Transgenderism 165 (2012)20
Florence Ashley, Gatekeeping Hormone Replacement Therapy for Transgender Patients Is Dehumanising, 45 J. Med. Ethics 480 (2019) 11
Hilary Cass, The Cass Review Independent Review of Gender Identity Services for Children and Young People: Interim Report (2022)11
Hilary Cass, The Cass Review Independent Review of Gender Identity Services for Children and Young People: Final Report (2024)19
Isabel Boyd et al., Care of Transgender Patients:  A General Practice Quality Improvement Approach, 10 Healthcare 11 (2022)12
James M. Cantor, Transgender and Gender Diverse Children and Adolescents: Fact-Checking of AAP Policy, J. Sex & Marital Therapy 307 (2019) 5, 10, 18
Jason Rafferty et al., Ensuring Comprehensive Care and Support for Transgender and Gender-Diverse Children and Adolescents, 142 Pediatrics 1 (2018)
Jay Cohn, <i>The Detransition Rate Is Unknown</i> , 52 Archives of Sex. Behav. 1937 (2023)
Jiska Ristori & Thomas D. Steensma, Gender  Dysphoria in Children, 28 Int. Rev. Psych. 1 (2016)

Julia Temple Newhook et al., A Critical Commentary on Follow-Up Studies and "Desistance" Theories about Transgender and Gender-Nonconforming Children, 19 Int'l J. Transgenderism 212 (2018) 13, 14, 15
Kenneth J. Zucker, Adolescents with Gender Dysphoria: Reflections on Some Contemporary Clinical and Research Issues, 48 Archives of Sex.Behav. 1 (2019)
Kenneth J. Zucker, <i>The Myth of Persistence</i> , 19 Int'l J. Transgenderism 231 (2018)14, 15
Kristina R. Olson et al., Gender Identity 5 Years  After Social Transition, 150 Pediatrics 1 (2022)
Kristina R. Olson, Prepubescent Transgender Children: What We Do and Do Not Know, 55 J. Am. Acad. Child & Adolescent Psych. 155 (2018)
Leor Sapir, Finland Takes Another Look at Youth Gender Medicine, Tablet, Feb. 21, 2023 19, 20
Lisa Littman, Rapid-Onset Gender Dysphoria in Adolescents and Young Adults: A Study of Parental Reports, 13 PLoS ONE 1 (2018)
Michael Biggs, The Dutch Protocol for Juvenile Transsexuals: Origins and Evidence, J. Sex & Marital Therapy 1 (2022)6
NHS England, Supporting Children and Young People with Gender-related Questions or Distress and Their Sexual Orientation, MindEd, July 11, 2023

Pablo Expósito-Campos and Roberto D'Angelo, Letter to the Editor: Regret after Gender-affirmation Surgery: A Systematic Review and Meta-analysis of Prevalence, 9 Plastic and Reconstructive Surgery—Global Open 3951 (Nov. 2021)
Peggy T. Cohen-Kettenis et al., The Treatment of Adolescent Transsexuals: Changing Insights, 5 J. Sexual Med. 1892 (2008)
Robin Respaut & Chad Terhune, Number of Transgender Children Seeking Treatment Surges in U.S., Reuters, Oct. 6, 2022
Ruth Hall et al., Access to Care and Frequency of Detransition among a Cohort Discharged by a UK National Adult Gender Identity Clinic: Retrospective Case-Note Review, 7 BJPsych Open 1 (2021)
Ruth Hall et al., Impact of Social Transition in Relation to Gender for Children and Adolescents: A Systematic Review, Archives Disease Childhood 1 (2024) 2, 18
Steward L. Adelson et al., Practice Parameter on Gay, Lesbian, or Bisexual Sexual Orientation, Gender Nonconformity, and Gender Discordance in Children and Adolescents, 51 J. Am. Acad. Child & Adolescent Psych. 957 (2012)6
Thomas D. Steensma & Peggy T. Cohen-Kettenis, Gender Transitioning Before Puberty?, 40 Archives of Sex. Behav. 649 (2011)
Thomas D. Steensma et al., Factors Associated with Desistence and Persistence of Childhood Gender Dysphoria: A Quantitative Follow-Up Study, 52 J. Am. Acad. Child Adolescent Psych. 582 (2013)

Valeria P. Bustos et al., Regret After Gender-
Affirmation Surgery: A Systematic Review and
Meta-Analysis of Prevalence,
9 Plastic and Reconstructive Surgery—
Global Open 3477 (2021)10
William Byne et al., Report of the APA Task Force on Treatment of Gender Identity Disorder 4 (2012)6
Wylie C. Hembree et al., Endocrine Treatment of
Gender-Dysphoric/ Gender-Incongruent Persons:
An Endocrine Society* Clinical Practice
Guideline, 102 J. Clinical Endocrinology &
Metabolism 3869 (2017)

#### INTEREST OF AMICI CURIAE<sup>1</sup>

The Manhattan Institute (MI) is a nonprofit policy research foundation whose mission is to develop and disseminate ideas that foster individual responsibility and agency across multiple dimensions. It has sponsored scholarship and filed briefs opposing regulations that interfere with constitutional liberties.

Leor Sapir, Ph.D., is a fellow at MI, where his research focuses on pediatric gender medicine and medical policy. His academic work, including his dissertation on Title IX, investigated how America's political culture and constitutional government shape public policy on matters of civil rights.

Amici file this brief because the right to direct the upbringing of one's children is among the most important liberties protected by the Constitution, one this Court has upheld repeatedly. Accordingly, amici believe that public-school teachers and staff, as state employees, are legally required to inform parents and respect parental direction regarding their children's health, including issues of gender and social transitioning. This brief highlights important medical research in that regard.

#### BACKGROUND AND SUMMARY OF ARGUMENT

Decades of research have consistently shown that most children with gender dysphoria (GD) and most clinically referred children with gender-variant behavior come to terms with their natal sex ("desist")

1

<sup>&</sup>lt;sup>1</sup> Rule 37 statement: Counsels of record for all parties were timely notified of this filing. No part of this brief was authored by any party's counsel, and no person or entity other than *amici* funded its preparation or submission.

by adulthood. Minors who are socially transitioned, however, are more likely to persist in their cross-gender feelings and, in time, seek medical interventions in the form of gonadotropin-releasing analogues (puberty blockers), cross-sex hormones, and surgeries. These interventions carry known and anticipated risks, including lifelong sterility, sexual dysfunction, mood disorders, and increased risk for cancer and heart disease.

In short, social transition is not a neutral act but an active intervention. But is it a beneficial one? The only systematic review of evidence on this question to date reported an "absence of robust evidence of the benefits or harms of social transition for children and adolescents." Ruth Hall et al., Impact of Social Transition in Relation to Gender for Children and Adolescents: A Systematic Review, Archives Disease Childhood 1, 1 (2024). A comprehensive assessment of over four decades of research does suggest, however, that social transition can lock in a temporary phase of development, leading to unnecessary identity medicalization and iatrogenic harm. Consequently, its use in schools falls squarely within parents' fundamental right to guide the healthcare decisions of their children.

By facilitating social transition without parental knowledge, the Eau Claire Area School District's Administrative Guidance for Gender Identity Support ("Gender Identity Support Policy") infringes on this fundamental right. Accordingly, the Court should grant certiorari to make clear to parents, schools, and courts across the nation that parents have standing to challenge school policies that withhold critical information from them about their children's mental health and well-being.

#### ARGUMENT

## I. Social Transition Constitutes a Mental-Health Intervention for Children Who Would Otherwise Likely Desist in Their Adopted Gender Identity before Adulthood

"Social transition" refers to the use of youths' preferred names and pronouns, access to sex-specific accommodations, and, in some cases, practices such as breast-binding and genital-tucking. Healthcare professionals worldwide have recognized transition as an active psychological intervention. Research strongly suggests that the vast majority of gender-dysphoric youths will naturally "desist," growing to feel comfortable with their natal sex. But social transition risks inhibiting this ordinary development, solidifying an otherwise passing phase of identity discordance past adolescence and, in turn, raising the potential for unnecessary medicalization.

In other words, social transition, far from reliving gender-related discomfort, may encourage these feelings to continue far longer than they would without it. Those who facilitate social transitions thus take part in a psychological intervention with potentially enormous ramifications for the well-being of children and adolescents.

#### A. Medical Research Worldwide Demonstrates That Social Transition Is a Mental-Health Intervention with Medical Implications for Children and Adolescents

The risks of early social transition were acknowledged by the Dutch clinicians who pioneered pediatric gender transition. In 2012, they recommended that young children not socially

transition before puberty on two grounds: (1) that most gender-dysphoric children will not persist in their adopted gender identity through adolescence; and (2) that such non-persisting youths should be prevented "from having to make a complex change back to the role of their natal gender," which research had suggested would be difficult. Annelou L. C. de Vries & Peggy T. Cohen-Kettenis, *Clinical Management of Gender Dysphoria in Children and Adolescents: The Dutch Approach*, 59 J. Homosexuality 301, 320 (2012).

The Dutch team also noted the danger of early social transition even for minors who do go on to full medical transition. Because medical transition cannot literally change a person's sex, they reasoned, it is important to ground the patient in reality and lower expectations about what drugs and surgeries can accomplish. The problem with "early transitions," they warned, "is that some children who have done so (sometimes as preschoolers) barely realize that they are of the other natal sex." *Id.* at 308. They develop a sense of reality so different from physical reality that acceptance of the protracted treatments they will later need is made unnecessarily difficult. Id. See also Thomas D. Steensma & Peggy T. Cohen-Kettenis, Gender Transitioning Before Puberty?, 40 Archives of Sex. Behav. 649, 649–50 (2011) (predicting "that the drawbacks of having to wait until early adolescence ... may be less serious than having to make a social transition twice").

Strikingly, in a 2008 article, the Dutch clinicians suggested that, given a "80-95%" desistence rate for gender dysphoria in children, a "real life test" or "real life experience" (*i.e.*, social transition) should be postponed until adolescence, and then only after an initial diagnosis of "gender identity disorder." Peggy T.

Cohen-Kettenis et al., The Treatment of Adolescent Transsexuals: Changing Insights, 5 J. Sexual Med. 1892, 1893 (2008). Social transition and pharmacological puberty suppression, they suggested, are both part of a prolonged diagnostic phase in the clinical management of youth gender dysphoria. Id. Both were to be used to discern the need for additional, more invasive, interventions—such as surgery. The Dutch thus seemed to have viewed social transition as less reversible than the administration of puberty blockers, an obvious medical intervention.

1. In most cases, childhood-onset gender dysphoria remits naturally by adulthood, but social transition may contribute to the persistence of gender dysphoria.

The Dutch researchers' cautious approach to social transition and their warnings about its risks are buttressed by <u>decades of research</u> finding that most children with gender identity issues come to terms with their natal sex, typically during adolescence. Those studies found <u>desistence rates of between 61 and 100 percent</u>, with specific percentages as follows in chronological order of publication: 75; 87.5; 100; 95.5; 90; 98; 87.5; 61; 88; 63; 87.7. James M. Cantor, *Transgender and Gender Diverse Children and Adolescents: Fact-Checking of AAP Policy*, J. Sex & Marital Therapy 307, 313 (2019) (collecting 11 studies from 1972 to 2019).

Of note, the studies found not only that most gender-dysphoric children eventually desist, but that a majority of natal males (63–100 percent) and a substantial minority of natal females (32–50 percent) who desisted later turned out to be gay or lesbian, not

transgender. Cross-gender feelings and behaviors in children are thus thought to be more predictive of later same-sex attraction than of lifelong gender dysphoria and trans identity. Early social transition may hinder healthy development of gender-nonconforming homosexual children. See also Michael Biggs, The Dutch Protocol for Juvenile Transsexuals: Origins and Evidence, J. Sex & Marital Therapy 1, 5 (2022).

The American Psychiatric Association observed in a 2012 literature review that "only a minority" of those diagnosed with childhood gender identity disorder "will identify as transsexual or transgender in adulthood (a phenomena termed persistence), while the majority will become comfortable with their natal gender over time (a phenomena called desistance)." William Byne et al., Report of the APA Task Force on Treatment of Gender Identity Disorder 4 (2012). That same year, the American Academy of Child and Adolescent Psychiatry acknowledged "longitudinal evidence that gender discordance persists in only a small minority of untreated cases arising in childhood," and warned that "further research is needed on predictors of persistence and desistence of childhood gender discordance as well as the long-term risks and benefits of intervention before any treatment to eliminate gender discordance can be endorsed." Steward L. Adelson et al., Practice Parameter on Gay, Lesbian, or Bisexual Sexual Orientation, Gender Nonconformity, and Gender Discordance in Children and Adolescents, 51 J. Am. Acad. Child & Adolescent Psych. 957, 968 (2012).

A major concern among researchers and clinicians who treat gender-diverse youth is that social transition will inhibit that natural remission and solidify an otherwise passing phase of identity discordance. For example, the Endocrine Society cautions that children who have socially transitioned "may have great difficulty in returning to the original gender role upon entering puberty," and that social transition "has been found to contribute to the likelihood of persistence." Wylie C. Hembree et al., Endocrine Treatment of Gender-Dysphoric/Gender-Incongruent Persons: An Endocrine Society Clinical Practice Guideline, 102 J. Clinical Endocrinology & Metabolism 3869, 3879 (2017).

One study found that childhood social transition was a factor associated with persistence. Thomas D. Steensma et al., Factors Associated with Desistence and Persistence of Childhood Gender Dysphoria: A Quantitative Follow-Up Study, 52 J. Am. Acad. Child Adolescent Psych. 582, 588 (2013). The most relevant question, however, pertains to the nature of that association. Are children more likely to be identified as transgender early on and then socially transitioned, or does social transition lock in cross-gender feelings and make it harder for gender-dysphoric kids to come to terms with their bodies?

A 2022 study, published in *Pediatrics*, challenges the conventional wisdom about desistance described above. Kristina R. Olson et al., *Gender Identity 5 Years After Social Transition*, 150 Pediatrics 1, 1 (2022) (special article). Based on their observation of 317 children, psychologist Kristina Olson and her colleagues claim to show that young children who are socially transitioned and "supported" in their new gender identity rarely change their minds. *Id.* at 6.

To be eligible for participation in the study, candidates had to have completed a full, "binary" (male-to-female or female-to-male) social transition.

Id. at 2. By the end of the study's five-year follow-up term, 3.5 percent of the children had replaced their male or female self-identification with a "non-binary" one, while 2.5 percent had "retransitioned" (i.e., come to terms with and learned to accept their natal sex). Id. at 3. For the study's authors and supporters of the "gender-affirmative" approach, this was good news: it confirmed the oft-repeated claim that "trans kids know who they are" and that children benefit from having adults agree with ("affirm") their asserted gender.

The serious problem with this interpretation is that it lacks "equipoise," which refers to the requirement that investigators show genuine uncertainty about an intervention's effects. See, e.g., Benjamin Freedman, Equipoise and the Ethics of Clinical Research, 317 New Eng. J. Med. 141, 141 (1987). Olson et al. failed to consider alternative explanations for why their findings conflicted with all previous research on persistence/desistence. With one partial exception, the children in the earlier studies did not undergo social transition and, with no exception, the studies yielded high rates of desistence. In Olson's study, however, all the children had been fully socially transitioned, and almost none desisted.

One interpretation of this discrepancy, favored by Olson and her colleagues, is that virtually all the children who participated in the 2022 study were "true transgender" children. See Olson, supra, at 4–6. But another explanation, overlooked by the authors, is that social transition itself caused them to persist, creating a self-fulfilling prophecy. Recall the Dutch clinicians' warnings that social transition can disrupt a child's grasp of reality and make coming to terms with his or her natal sex more difficult. See de Vries & Cohen-Kettenis, supra, at 308.

Despite its authors' interpretation, the Olson study suggests that social transition may in fact be a powerful psychological intervention with potential to lock in gender incongruence. If true, the consequences are serious: at least 60 percent of the children in the study had commenced hormonal interventions, which carry significant health risks, at the five-year follow-up. Olson, supra, at 2. If some of these children might have desisted and avoided unnecessary medicalization, then their social transition was the cause of iatrogenic harm.

# 2. Transgender identity in adolescents is also likely unstable.

Proponents of social and medical gender reassignment for minors argue that when gender dysphoria begins in childhood and intensifies at the outset of puberty, the chances of desistence are very slim. This belief is not supported by evidence.

First, in the 11 desistence studies discussed above. some of the minors who desisted did so after they had entered adolescence. Cantor, supra, at 5. That's why the Endocrine Society's guidelines mention that GD/gender incongruence "childhood invariably persist into adolescence and adulthood." Hembree et al., supra, at 3876 (emphasis added). Second, gender clinics in a variety of countries and researchers who study gender dysphoria in youth have observed a new patient cohort that does not fit the profile of the youth who participated in the original Dutch study and for whom the Dutch pioneered pediatric gender transition. See E. Abruzzese et al., The Myth of "Reliable Research" in Pediatric Gender Medicine: A Critical Evaluation of the Dutch Studies and Research that Has Followed, J. Sex & Marital Therapy 1, 12–13 (2023). This new cohort of minors, which accounts for most of the meteoric increase in the number of minors seeking gender transition services over the past decade, is comprised of young people who did not have gender-identity issues in childhood and whose gender-dysphoric symptoms began, suddenly, after the start of puberty. *Id.* Most are natal girls with comorbid mental-health problems. Id. The very fact that these teenagers exist suggests that transgender identity is neither innate nor immutable. See also Kenneth J. Zucker, Adolescents with Gender Dysphoria: Reflections on Some Contemporary Clinical and Research Issues, 48 Archives of Sex. Behav. 1, 7 (2019); Lisa Littman, Rapid-Onset Gender Dysphoria in Adolescents and Young Adults: A Study of Parental Reports, 13 PLoS ONE 1, 30–33 (2018).

Third, researchers are increasingly acknowledging the phenomenon of regret and detransition. Claims about regret and detransition being extremely rare less than 2 percent, by some accounts—are based on studies done mainly on adults who transitioned as adults. See Valeria P. Bustos et al., Regret After Gender-Affirmation Surgery: A Systematic Review and Meta-Analysis ofPrevalence. 9 Reconstructive Surgery—Global Open 3477, (Mar. 2021). These studies suffer from high drop-out rates, short follow-up times, and highly restrictive definitions of detransition. Pablo Expósito-Campos and Roberto D'Angelo, Letter to the Editor: Regret after Gender-affirmation Surgery: A Systematic Review and Meta-analysis ofPrevalence, 9 PlasticReconstructive Surgery—Global Open 3951 (Nov. 2021). The very few adolescents included in these statistics were all transitioned under the Dutch protocol, a relatively conservative approach that contrasts with the affirmative approach practiced in American clinics. See Abruzzese et al., supra, at 14–15. It is irresponsible to say that the extremely low rates of regret/detransition observed in earlier studies apply to the majority of minors seeking social or medical gender transition today. These are distinct clinical cohorts with different presentations and clinical needs, and there is no high-quality research on the adolescent-onset group.

Unlike the more conservative Dutch protocol, which requires a childhood diagnosis of gender dysphoria that intensifies in adolescence and no serious psychological comorbidities, the affirmative approach regards adolescent-onset gender dysphoria—even when it appears abruptly and develops rapidly—as a valid transgender identity and considers co-occurring mental health problems as secondary to gender identity problems.

In her interim report to the U.K.'s National Health Service, Dr. Hilary Cass called this problem "diagnostic overshadowing": once the clinician identifies gender as a source of distress, all other problems, including ones that might be causing the gender distress, are ignored. Hilary Cass, The Cass Review Independent Review of Gender Identity Services for Children and Young People: Interim Report 17 (2022). Some prominent proponents of the gender-affirmative model for youth in the United States have argued that there should be "gatekeeping" at all, only "informed consent." See, e.g., Florence Ashley, Gatekeeping Hormone Replacement Therapy for Transgender Patients Is Dehumanising, 45 J. Med. Ethics 480, 480-81 (2019). Lowering the thresholds for medical treatment is likely to increase the rate of false positives and thus the rate of regret.

Recent studies have shown high rates of regret and detransition. A study using data from the U.S. military healthcare system found that 30 percent of those who started treatments discontinued them within four vears. See Christina M. Roberts et al., Continuation of Gender-Affirming Hormones Among Transgender Adolescents and Adults, 107 J. Clinical Endocrinology & Metabolism 3937, 3937 (2022). Another study from the U.K. in 2021 found that 10 percent of those treated at an adult transgender clinic detransitioned within 16 months of receiving treatment. Ruth Hall et al., Access to Care and Frequency of Detransition among a Cohort Discharged by a UK National Adult Gender Identity Clinic: Retrospective Case-Note Review, 7 BJPsych Open 1, 7 (2021). An additional 22 percent disengaged from the clinic before completing their treatment. Id. at 5. Another study on adults found a rate of regret or detransition of 12 percent and a rate of discontinuation of 20 percent. Isabel Boyd et al., Care of Transgender Patients: A General Practice Quality Improvement Approach, 10 Healthcare 11 (2022). The authors of the study noted that "the detransition rate found in this population is novel and questions may be raised about the phenomenon of overdiagnosis, overtreatment, or iatrogenic harm as found in other medical fields." Id. at 13.

In sum, today's evidence suggests that transgender identity is less stable in adolescents, or even adults, than social-transition advocates assert. The true rate of regret and detransition is not known. See generally Jay Cohn, The Detransition Rate Is Unknown, 52 Archives of Sex. Behav. 1937 (2023). High-quality research is necessary, especially on teenagers with complex presentations of gender-related distress, to know how many will experience regret or detransition.

#### B. Objections to the Persistence and Desistence Literature Do Not Withstand Scrutiny

To interpret away the overwhelming evidence that childhood gender dysphoria typically remits by adulthood, some researchers and transgender activists have raised methodological objections to the studies discussed above. These all fail upon close scrutiny. One such objection is that many or most of the children in the desistence studies were not "truly transgender." As a result, the objection asserts, this desistence finding is artificially inflated. See, e.g., Julia Temple Newhook et al., A Critical Commentary on Follow-Up Studies and "Desistance" Theories about Transgender and Gender-Nonconforming Children, 19 Transgenderism 212, 214-16 (2018); Kristina R. Olson, Prepubescent Transgender Children: What We Do and Do Not Know, 55 J. Am. Acad. Child & Adolescent Psych. 155, 155–56 (2016). This argument is built on two claims. First, earlier studies inappropriately included not just children diagnosed with gender-identity disorder—or GID, as it was called at the time—but also children who were merely gender-nonconforming. Newhook et al., supra, at 215-16. Second, even the children who were diagnosed with GID were not necessarily transgender because the diagnostic criteria under the American Psychiatric Association's Diagnostic and Statistical Manual of Mental Disorders (DSM) Edition III (1980), III-R (1987), and IV (1994) were not as demanding as those for gender dysphoria in DSM-5 (2013). *Id*.

The first claim fails because subsequent research has borne out the durability of the desistence literature. Dr. Kenneth Zucker, an internationally renowned expert in pediatric gender dysphoria who chaired the DSM-5's Work Group on Sexual and Gender Identity Disorders and helped write the diagnostic guidelines for "gender identity disorder" in the DSM-III-R and DSM-IV, conducted a re-analysis of the earlier studies in the light of criticism about the validity of their findings. See Kenneth J. Zucker, The Myth of Persistence, 19 Int'l J. Transgenderism 231, 233–34 (2018). He divided the children in these studies into two groups: those who did not meet the thresholds for GID but were still dysphoric enough to require referral to a specialized gender clinic, and those who did meet the diagnostic threshold. Those who were subthreshold desisted at a rate of over 90 percent, while those who met the threshold desisted at a rate of almost 70 percent. Id. at 234–35.

The second claim does not fare better. The DSM-5 includes two criteria for diagnosing gender dysphoria in children: at least six months of "marked incongruence between one's experienced/expressed gender and assigned gender" and "clinically significant distress or impairment in social, school, or other important areas of functioning." Am. Psych. Ass'n, DSM-5, at 452.

The DSM-5's innovation on previous editions is its inclusion, within the first of these two criteria ("marked incongruence"), of sub-criterion A1: "A strong desire to be of the other gender or insistence that one is the other gender (or some alternative gender different from one's assigned gender)." *Id.* Critics of the desistence literature argue that A1 represents a raising of the diagnostic threshold, and consequently, many or most of the children diagnosed with GID under previous DSM editions would not have met the DSM-5's threshold. Newhook et al., *supra*, at 214–15.

But the addition of A1 is of no real clinical significance, and the objection is misguided. The diagnostic criteria from the DSM-IV include "a strong and persistent cross-gender identification" and, in children, a "[r]epeatedly stated desire to be, or insistence that he or she is, the other sex." Am. Psych. Ass'n, DSM-IV, at 581 (DSM-IV-TR: Gender Identity Disorder in Children (302.6) and Gender Identity Disorder in Adolescents or Adults (302.85)). And the criticism largely ignores the crucially important guideline accompanying the text of the diagnostic criteria, advising that the "disorder is not meant to describe a child's nonconformity to stereotypic sex-role behavior" and that it instead "represents a profound disturbance of the individual's sense of identity with regard to maleness or femaleness." Id. at 580. It is thus wrong to infer that the DSM-IV included mere gendernonconformity in its description of gender identity disorder.

An analysis of the differences between the DSM-IV and DSM-5 by the Dutch team that pioneered pediatric gender transition concludes that "both editions ... of gender-identity related diagnoses seem reliable and convenient for clinical use." Annelou L. C. de Vries et al., Reliability and Clinical Utility of Gender Identity-Related Diagnoses: Comparisons Between the ICD-11, ICD-10, DSM-IV, and DSM-5, 8 LGBT Health 133, 133 (2021). Dr. Zucker, who helped write the diagnostic guidelines for both DSM editions, has written: "It is my clinical opinion that the similarities across the various iterations of the DSM are far greater than the differences." Zucker, The Myth of Persistence, supra, at 234.

It should finally be noted that the assumption that the DSM-5's diagnostic threshold is higher than that of earlier editions is itself dubious. The number of minors receiving a gender dysphoria diagnosis has skyrocketed since the publication of the DSM-5. In the United States, according to data collected by Komodo Health and reported by Reuters, diagnoses in youth ages 6-17 rose by approximately 20 percent annually between 2017 and 2020, and by approximately 70 percent between 2020 and 2021, for a total of 121,882 new diagnoses added during these years. Robin Respaut & Chad Terhune, Number of Transgender Children Seeking Treatment Surges in U.S., Reuters, Oct. 6, 2022. Since these data are based on insurance claims, they likely represent an undercount of the true number. It seems that the DSM-5 makes it easier, not harder, to receive a diagnosis. In short, there is no basis to believe that four decades of research have yielded an artificially inflated rate of desistence.

#### C. Clinicians Have Not Demonstrated a Consistent Ability to Distinguish between Transgender and Gender-Nonconforming Youths

Some supporters of social transition argue that clinicians can reliably distinguish persisters from desisters in childhood. Children who express gender identity in a way that is "insistent, persistent, and consistent" (IPC), these supporters argue, can be regarded as "true transgender" children. The ability to avoid false positives means that clinicians can recommend social transition even if many or most children desist, and even if social transition is inappropriate for children who appear to be, but are not, transgender.

Proponents of IPC point to research showing that some factors—including age, natal sex, and diagnosis

of GID/GD—are associated with a higher rate of persistence. They argue that children in whom one or more of these factors appear are "true transgender." The problem with this argument is that it tries to infer *individual* predictions from *population* data. As one group of experts explains:

Factors predictive for the persistence of GD have been identified on a group level, with higher intensity of GD in childhood identified as the strongest predictor for a future gender dysphoric outcome. The predictive value of the identified factors for persistence are, however, on an individual level less clear cut, and the clinical utility of currently identified factors is low.

Jiska Ristori & Thomas D. Steensma, Gender Dysphoria in Children, 28 Int. Rev. Psych. 1, 6 (2016) (internal citations omitted). The Endocrine Society's 2017 guidelines on treatment of gender dysphoric youth likewise recognize that "[w]ith current knowledge, we cannot predict the psychosexual outcome for any specific child." Hembree et al., supra, at 3876.

#### D. U.S.-Based Medical Groups Are Out of Step with World Health Authorities' Recognition of the Risks of Pediatric Social Transition

The American Academy of Pediatrics has called for automatic gender affirmation (social transition) of minors, irrespective of their age, since 2018. Jason Rafferty et al., Ensuring Comprehensive Care and Support for Transgender and Gender-Diverse Children and Adolescents, 142 Pediatrics 1, 57 (2018). The AAP

statement has been subjected to thorough criticism for its inaccuracies and misrepresentations, see Cantor, supra, at 313, and contrasts sharply with medical authorities abroad. Even leading members of the World Professional Association for Transgender Health have admitted that the AAP's statement has "a very weak methodology." Appendix A to Supplemental Expert Report of James Cantor, Ph.D., at iv, Boe v. Marshall, No. 2:22-cv-00184-LCB-CWB (M.D. Ala. June 24, 2024) (quoting internal WPATH email).

In 2023, the NHS published a module for schools in which it reiterated that social transition is "a complex and should be considered an intervention." NHS England, Supporting Children and Young People with Gender-related Questions or Distress and Their Sexual Orientation, MindEd, July 11, 2023, https://bit.ly/478qu8z. "Supporting a social transition without the involvement of parents or carers," the NHS emphasized, "can create complex difficulties within families and is not recommended. Secrets between parents or carers and their children are problematic and are likely to create further issues in the future." *Id.* As for the concern about abusive parents, "if there are significant concerns ... schools should seek careful and detailed safeguarding oversight to assess risks." Id.

The Cass Review, a comprehensive report on youth gender transition commissioned by the National Health Service of England and released in April 2024, commissioned a systematic review of the evidence for mental health outcomes of social transition in minors and found an "absence of robust evidence of the benefits or harms of social transition for children and adolescents." Hall, *Impact of Social Transition*, *supra*, at 1. The Review emphasized that social transition

should be thought of as an "active intervention because it may have significant effects on the child or young person in terms of their psychological functioning and longer-term outcomes." Hilary Cass, The Cass Review Independent Review of Gender Identity Services for Children and Young People: Final Report 158 (2024). For young children in particular, full social transition may lock in a temporary phase of identity confusion, making persistence and medicalization more likely. "Clinical involvement in the decision-making process should include advising on the risks and benefits of transition planned social as a intervention. referencing best available evidence. This is not a role that can be taken by staff without appropriate clinical training." Id. at 164. And crucially, for both children adolescents. "Outcomes for children adolescents are best if they are in a supportive relationship with their family. For this reason parents should be actively involved in decision making unless there are strong grounds to believe that this may put the child or young person at risk." *Id*.

Dr. Riittakerttu Kaltiala, chief psychiatrist at Tampere University's pediatric gender clinic in Finland, recently confirmed that "four out of five" children with gender dysphoria desist by adulthood. Leor Sapir, Finland Takes Another Look at Youth Medicine, Tablet. Feb. 21, Gender 2023. https://bit.ly/3YVwxZp. Asked to comment on a proposed law that would grant minors the ability to define their gender for purposes of government documents, Dr. Kaltiala said that while it is "important to accept [children] as they are," "negating the body" by confirming that a child's gender selfperception is real can send the child "a message that there is something wrong with him or her." Annika Mutanen, A Professor Who Treats Adolescent Gender Anxiety Says No to Minors' Legal Gender Correction, Helsingin Sanomat, Jan. 27, 2023 (translated from Finnish using Google Translate). The Finnish Paediatric Society and Finnish Medical Association both objected to legal gender self-identification for minors, and the proposal was defeated. See Sapir, supra.

Even U.S.-based World the Professional Association for Transgender Health (WPATH) recognized that "[t]he current evidence base is insufficient to predict the long-term outcomes of completing a gender role transition," and that the desire for social transition "may reflect an expression of their gender identity" but it could also "be motivated by other forces." Because "[a] change back to the original gender role can be highly distressing," parents should rely on "[m]ental health professionals" to help them "make decisions regarding the timing and process of any gender role changes for their young children." Eli Coleman et al., Standards of Care for the Health of Transsexual, Transgender, and Gender Nonconforming People, Version 7, 13 Int'l J. Transgenderism 165, 176 (2012).

WPATH's Version 8 now recommends social transition for children but only "when it would be beneficial." Eli Coleman et al., Standards of Care for the Health of Transgender and Gender Diverse People, Version 8, 23 Int'l J. Transgender Health S1, S76 (Suppl. 1) (2022). Who determines that? According to WPATH, "health care professionals [should] discuss the potential benefits and risks of a social transition with families who are considering it." Id. at S69, S77. For adolescents, WPATH recommends that "family members ... work collaboratively" with "community

members" such as school officials, "unless [families'] involvement is considered harmful to the adolescent." *Id.* at S52. WPATH's new recommendations are not based on systematic reviews of evidence and discuss only the benefits of social transition, yet even they emphasize the need for clinical consultation.

#### II. U.S. Courts Have Recognized Social Transition as a Medical Intervention

Lawsuits over the past decade have raised the question of how schools must accommodate transgender-identified students under Title IX of the 1972 Education Amendments, 20 U.S.C. §§ 1681-1688, and the Equal Protection Clause, U.S. Const. amend. XIV § 1. Transgender-identified plaintiffs and their counsel have argued, and in most cases, courts have agreed, that using restrooms and other sexspecific facilities are a crucial part of a treatment protocol for a "medically diagnosed and documented condition." Whitaker v. Kenosha Unified Sch., Dist., 858 F.3d 1034, 1050 (7th Cir. 2017).

A therapeutic rationale for social transition and gender identity-based access to restrooms in schools has been stated by transgender-identified students and their representatives in multiple circuits. See, e.g., Br. of Amici Curiae Am. Acad. Pediatrics in Support of Defendants-Appellees and Affirmance at 6, Soule v. Ct. Ass'n of Schs., No. 3:20-cv-00201 (2nd Cir. 2022); Br. of Amici Curiae Am. Acad. of Pediatrics at 14, Doe v. Boyertown Area Sch. Dist., 897 F.3d 518 (3d Cir. 2018); Br. of Amici Curiae Med., Pub. Health, and Mental Health Orgs. in Support of Plaintiff-Appellee at 3, Grimm v. Gloucester Cnty. Sch. Bd., 972 F.3d 586 (4th Cir. 2020); Complaint at 21–24, Bd. of Educ. of the Highland Local Sch. Dist. v. U.S. Dep't of Educ., 208

F. Supp. 3d 850 (S.D. Ohio 2016); Whitaker v. Kenosha *Unified Sch. Dist.*, 858 F.3d 1034, 1050 (7th Cir. 2017); Br. of Amici Curiae Am. Acad. of Pediatrics at 3–4, Parents for Privacy v. Dallas Sch. Dist. No. 2, 949 F.3d 1210 (9th Cir. 2020); Adams v. Sch. Bd. of St. Johns Cnty., No. 18-13592, at \*7–9 (11th Cir. 2022) (Pryor, J., dissenting). Five circuit courts agreed that social transition and, by extension, use of accommodations designated for the sex students identify with is important for the rapeutic treatment. See Doe v. Boyertown Area Sch. Dist., 897 F.3d 518, 529-31 (3d Cir. 2018); Grimm v. Gloucester Cnty. Sch. Bd., 972 F.3d 586, 615, 619 (4th Cir. 2020); Dodds v. U.S. Dep't of Educ., 845 F.3d 217, 221-22 (6th Cir. 2016); Whitaker v. Kenosha Unified Sch. Dist., 858 F.3d 1034, 1050 (7th Cir. 2017); Parents for Privacy v. Dallas Sch. Dist. No. 2, 949 F.3d 1210, 1210 (9th Cir. 2020).

## III. The Eau Claire Area School District's Gender Identity Support Policy Prevents Parents from Assessing the Benefits and Risks of Social Transition and Other Mental-Health Aspects of Gender Care

When social transition is properly understood as an active mental-health intervention, it becomes clear that the Eau Claire Area School District's Gender Identity Support Policy infringes on the parental rights of Petitioner's members, causing injury.

Because social transition is a powerful mental-health intervention, it is a decision that should be made by, or in consultation with, the parents or legal guardians of students. This decision falls squarely within the ambit of parents' fundamental right to make healthcare decisions for their children. See, e.g., Stanley v. Illinois, 405 U.S. 645, 651 (1972) (affirming

parents' rights "in the companionship, care, custody, and management" of their children); Lassiter v. Dep't of Soc. Servs., 452 U.S. 18, 27 (1981) (quoting Stanley and reaffirming that parents' interest over their children "undeniably warrants deference and, absent a powerful countervailing interest, protection").

This Court has been clear and consistent in stressing that parents or legal guardians are the ones best positioned to understand and address the needs of children and adolescents, including unique mental health needs. See, e.g., Troxel v. Granville, 530 U.S. 57, 66–69 (2000). Overriding their discretion requires an exceedingly persuasive justification, compelling evidence that physical or emotional abuse will ensue if a student's desire for social transition is disclosed. See Santosky v. Kramer, 455 U.S. 745, 752-54 (1982). Even in these very limited circumstances, school staff should not be at liberty to perform social transitions but should instead consult the appropriate state child welfare services.

By facilitating the active psychological intervention of social transition without parental knowledge, the Gender Identity Support Policy prevents parents from exercising their fundamental right to determine the healthcare decisions of their children. See Parham v. J.R., 442 U.S. 584, 604 (1979); Colon v. Collazo, 729 F.2d 32, 34 (1st Cir. 1984). That right prevails even when children or school officials dislike or disagree with fit parents' exercise of their discretion. This Court has consistently protected the "cardinal" principle that "that the custody, care and nurture of the child reside first in the parents" and "respected the private realm of family life which the state cannot enter." Prince v. Massachusetts, 321 U.S. 158, 166 (1943).

Given the well-established risks associated with social transition, children should not be exposed to the potential for medicalization and iatrogenic harm before parents have standing to sue. Contrary to the Seventh Circuit's opinion, Petitioner's injury is not grounded in a "worry that may never come to pass," App. 2, but necessarily follows from the Gender Identity Support Policy's option to keep parents unaware of their child's gender status. See App. 66, 72–73.

When school staff make decisions about social transition in secret, they usurp the authority of parents and violate their well-established constitutional rights. Most importantly, they put the mental and physical health of minors at risk.

#### CONCLUSION

The Court should grant this petition.

Respectfully submitted,

Ilya Shapiro

Counsel of Record

MANHATTAN INSTITUTE
52 Vanderbilt Ave.

New York, NY 10017
(212) 599-7000
ishapiro@manhattan.institute

July 8, 2024